

## *The California Citrus Variety Improvement Program*

**I**N A PREVIOUS REPORT (3), the historical background, objectives, facilities and program for establishing and maintaining virus-free citrus stocks in California were outlined. The program is a coordinated undertaking of the University of California Citrus Experiment Station, the California State Department of Agriculture, and the U.S. Department of Agriculture. The University is developing and will maintain primary sources ("foundation blocks") of healthy, true-to-name propagating material of improved citrus scion varieties and rootstocks. The University's phase is administered by the Citrus Variety Improvement Committee. The State Department of Agriculture will handle the regulatory and certification phase, the details of which have been outlined by Hiltabrand (1). The role of the U.S. Department of Agriculture in the importation of foreign varieties is outlined in a previous report (4).

This paper presents some data on the incidence of certain virus diseases in citrus selections of both domestic and foreign origin which have been indexed in connection with this program.

### *Domestic Indexing Program*

Details of the indexing procedures used are essentially those described by Wallace and Drake (4). Individual trees of varieties or strains of domestic (California) origin to be indexed are called "candidate" trees, and are selected by the Citrus Variety Improvement Committee. Normally, such candidate trees are mature, bearing trees which are known to be true to name or type, free of bud mutations, and apparently vigorous

per cent, of the foreign clones contained 1 or more of the 4 viruses listed above. Of the 30 that were positive for psorosis, 13 were discarded but 17 are being held for observation of fruit characters, and possible production of nucellar clones from any that are particularly desirable. It is evident that these viruses have a wide geographic distribution, since these accessions came from 25 different countries. The discovery of tristeza virus in introductions from 3 countries was the first evidence of the presence of tristeza in those countries. The data in Table 1 also indicate that these viruses are common in a wide range of species and varieties in the genus *Citrus* and that they may occur in certain citrus relatives.

Of the 57 accessions surviving the short-term indexing, 17 have completed the 3-year scion-stock indexing (4) carried on in the quarantine greenhouse. Of the 17, 5 lemon varieties, Berna, Ricote, La Porto, Mesero Jhamberi, 2 mandarin varieties, Clementine de Alcazar and China orange, and 1 orange, Sanguinello Moscato Cuscuna, have been moved to the Lindcove foundation block; others will be moved next spring.

The incidence of virus diseases in foreign accessions so far disclosed by short-term indexing is rather high. If the incidence of xyloporosis and exocortis turned up by the long-term indexing procedures to be carried out at Lindcove is as high as found by Norman (2) in Florida, perhaps fewer than 20 of the 57 introductions will prove to be free of detectable virus diseases. This rate of loss is extremely high for the time, effort, and expense involved and emphasizes the critical need for faster indexing procedures and dependable, simple ways of eliminating specific virus diseases from citrus tissues. A reasonably rapid means of exchanging new or untried citrus varieties and breeding material among workers in the United States and other citrus-growing areas of the world without exchanging diseases or pests, especially viruses, is essential if reasonable progress is to be made in improving citrus varieties through breeding and selection.

#### *Literature Cited*

1. HILTABRAND, W. F. 1959. Certification program for maintenance of virus-free propagation sources of citrus in California, p. 229-231. *In* J. M. Wallace [ed.], *Citrus Virus Diseases*. Univ. Calif. Div. Agr. Sci., Berkeley.
2. NORMAN, G. G. 1959. Florida State Plant Board programs for virus-free budwood, p. 237-242. *In* J. M. Wallace [ed.], *Citrus Virus Diseases*. Univ. Calif. Div. Agr. Sci., Berkeley.

PROCEEDINGS of the IOCV

TABLE 1. INCIDENCE OF CERTAIN VIRUS DISEASES IN CITRUS CANDIDATE TREES INDEXED IN THE CALIFORNIA CITRUS VARIETY IMPROVEMENT PROGRAM

Group or species	Accessions positive for				accessions negative
	tristeza	seedling yellows	psorosis	vein enation	
<i>Domestic accessions</i>					
Orange - Navel	3	—	1	2	11
Orange - Valencia	1	—	0	0	5
Orange - misc.	1	—	0	1	0
Eureka lemon	0	—	0	1	5
Lisbon lemon	0	—	0	2	7
Other lemon	0	—	2	0	2
Grapefruit	0	—	0	0	4
Mandarin	0	—	0	1	15
Lime	0	—	0	0	1
Tangelo	0	—	0	0	2
Hybrid - misc.	0	—	0	0	8
Totals, Domestic	5	—	3	7	60
<i>Foreign accessions</i>					
<i>C. aurantifolia</i>	0	0	0	1	3
<i>C. bergamia</i>	1	0	0	0	1
<i>C. grandis</i>	1	0	0	0	6
<i>C. jambheri</i>	2	0	0	0	1
<i>C. limon</i> <sup>a</sup>	3	1	6	0	9
<i>C. medica</i>	2	1	2	0	1
<i>C. natsudaikai</i> <sup>a</sup>	1	0	1	0	0
<i>C. nobilis, unshiu</i>	0	0	0	0	3
<i>C. paradisi</i>	0	0	0	0	1
<i>C. reticulata</i> <sup>a</sup>	1	0	2	0	7
<i>C. sinensis</i> <sup>a</sup>	0	1	18	0	12
<i>C. spp.</i>	6	2 <sup>b</sup>	0	1 <sup>b</sup>	10
<i>Fortunella japonica</i>	1 <sup>b</sup>	0	1 <sup>b</sup>	0	0
<i>Paramignya monophylla</i>	0	0	0	0	1
<i>Micromelum spp.</i> <sup>c</sup>	0	0	0	0	1
<i>Poncirus, spp.</i>	0	0	0	0	1
Totals, Foreign	18	5	30	2	57

<sup>a</sup>In a few of the earliest accessions, no further indexing was performed if clear-cut psorosis symptoms were present on the plant when received at Riverside. Falling in this category were 8 *C. sinensis*, 1 each of *C. limon*, *C. reticulata*, and *C. natsudaikai*. Had the regular indexing program been followed on these plants also, a few more accessions with more than 1 virus might have been recorded.

<sup>b</sup>One plant contained 2 viruses.

<sup>c</sup>Identity questionable.

28 have been moved to the foundation block at the University of California experimental citrus orchard at Lindcove in Tulare County. This is in an important citrus-growing area in the San Joaquin Valley located about 250 miles from Riverside, and is well isolated from other citrus-growing areas by mountains and arid, non agricultural regions. Recent surveys have detected neither tristeza nor vein enation in the area.

Because tristeza is not present in all citrus areas in California, quarantine regulations established by the California State Department of Agriculture prohibit the movement of citrus budwood or nursery trees from infested areas to noninfested areas. However, by special permit and under rigorous inspection by California State Department of Agriculture personnel, tristeza-free source trees produced and held in the University of California quarantine houses at Riverside, a tristeza-infested area, are permitted to move to the Lindcove site, a tristeza-free area. Many valuable citrus clones, including a number of nucellar origin, are represented in the Citrus Experiment Station variety collection at Riverside. The threat of infestation with tristeza and other insect-transmitted viruses present in the area lends urgency to moving virus-free propagative material of such clones to a tristeza-free location, such as the Lindcove foundation plot.

### *Foreign Indexing Program*

The introduction of fungus or bacterial diseases and insect or nematode pests on foreign citrus introductions is avoided by first propagating them in the greenhouses of the U.S. Department of Agriculture at Glenn Dale, Maryland. After about 2 years of careful inspection, clean material is sent to California for rigorous virus disease indexing, which involves separate greenhouse and screenhouse facilities isolated from those used for the indexing of domestic selections. The short-term indexing procedures are essentially the same as for the domestic selections, but the long-term indexing procedures are more rigorous and exacting (4).

In the 7 years since the foreign citrus budwood importation program (4) was started, 143 accessions have been received at Riverside. Of these, 13 were discarded because the variety or species was already established at Riverside, or because of insufficient descriptive information. Currently 20 accessions are being indexed for psorosis, tristeza, seedling yellows, and vein enation. Indexing for these viruses has been completed on 110 foreign citrus accessions. A summary is presented in Table 1.

In contrast with the domestic indexing results, 53 out of 110, or 48

per cent, of the foreign clones contained 1 or more of the 4 viruses listed above. Of the 30 that were positive for psorosis, 13 were discarded but 17 are being held for observation of fruit characters, and possible production of nucellar clones from any that are particularly desirable. It is evident that these viruses have a wide geographic distribution, since these accessions came from 25 different countries. The discovery of tristeza virus in introductions from 3 countries was the first evidence of the presence of tristeza in those countries. The data in Table 1 also indicate that these viruses are common in a wide range of species and varieties in the genus *Citrus* and that they may occur in certain citrus relatives.

Of the 57 accessions surviving the short-term indexing, 17 have completed the 3-year scion-stock indexing (4) carried on in the quarantine greenhouse. Of the 17, 5 lemon varieties, Berna, Ricote, La Porto, Mesero Jhamberi, 2 mandarin varieties, Clementine de Alcazar and China orange, and 1 orange, Sanguinello Moscato Cuscuna, have been moved to the Lindcove foundation block; others will be moved next spring.

The incidence of virus diseases in foreign accessions so far disclosed by short-term indexing is rather high. If the incidence of xyloporosis and exocortis turned up by the long-term indexing procedures to be carried out at Lindcove is as high as found by Norman (2) in Florida, perhaps fewer than 20 of the 57 introductions will prove to be free of detectable virus diseases. This rate of loss is extremely high for the time, effort, and expense involved and emphasizes the critical need for faster indexing procedures and dependable, simple ways of eliminating specific virus diseases from citrus tissues. A reasonably rapid means of exchanging new or untried citrus varieties and breeding material among workers in the United States and other citrus-growing areas of the world without exchanging diseases or pests, especially viruses, is essential if reasonable progress is to be made in improving citrus varieties through breeding and selection.

#### *Literature Cited*

1. HILTABRAND, W. F. 1959. Certification program for maintenance of virus-free propagation sources of citrus in California, p. 229-231. *In* J. M. Wallace [ed.], *Citrus Virus Diseases*. Univ. Calif. Div. Agr. Sci., Berkeley.
2. NORMAN, G. G. 1959. Florida State Plant Board programs for virus-free budwood, p. 237-242. *In* J. M. Wallace [ed.], *Citrus Virus Diseases*. Univ. Calif. Div. Agr. Sci., Berkeley.

REUTHER

3. REUTHER, W. 1959. A program for establishing and maintaining virus-free citrus stock, p. 215-217. *In* J. M. Wallace [ed.], Citrus Virus Diseases. Univ. Calif. Div. Agr. Sci., Berkeley.
4. WALLACE, J. M., and R. J. DRAKE. 1959. An indexing program to avoid viruses in citrus introduced into the United States, p. 209-214. *In* J. M. Wallace [ed.], Citrus Virus Diseases. Univ. Calif. Div. Agr. Sci., Berkeley.

*This is a report of the Citrus Variety Improvement Committee, which consists of three members of the Department of Horticulture: Dr. W. P. Bitters, Dr. J. W. Cameron, and Dr. Walter Reuther, Chairman; three members from the Department of Plant Pathology: Dr. E. C. Calavan, Dr. J. M. Wallace, and Dr. L. G. Weathers; and one representative of the U.S. Department of Agriculture: Dr. J. R. Furr. Special recognition is due Mr. E. M. Nauert, Assistant Specialist, and Mr. C. N. Roistacher, Laboratory Technician IV, for their very capable services in handling the actual details of the plant care, propagation, indexing, and record-keeping procedures.*